

Bounds on graph eigenvalues II

Vladimir Nikiforov

Department of Mathematical Sciences, University of Memphis,
Memphis TN 38152, USA, e-mail: *vnikifrv@memphis.edu*

February 2, 2008

Abstract

We prove three results about the spectral radius $\mu(G)$ of a graph G :

(a) Let $T_r(n)$ be the r -partite Turán graph of order n . If G is a K_{r+1} -free graph of order n , then

$$\mu(G) < \mu(T_r(n))$$

unless $G = T_r(n)$.

(b) For most irregular graphs G of order n and size m ,

$$\mu(G) - 2m/n > 1/(2m + 2n).$$

(c) Let $0 \leq k \leq l$. If G is a graph of order n with no $K_2 + \overline{K}_{k+1}$ and no $K_{2,l+1}$, then

$$\mu(G) \leq \min \left\{ \Delta(G), \left(k - l + 1 + \sqrt{(k - l + 1)^2 + 4l(n - 1)} \right) / 2 \right\}.$$

Keywords: *clique number, spectral radius, Turán graph, maximum degree, books*

1 Introduction

Our notation is standard (e.g., see [1]); specifically, we write $G(n)$ for a graph of order n and $\mu(G)$ for the maximum eigenvalue of the adjacency matrix of G .

Write $T_r(n)$ for the r -partite Turán graph of order n and let $G = G(n)$. In [7] it is shown that if G is r -partite, then $\mu(G) < \mu(T_r(n))$ unless $G = T_r(n)$. On the other hand, Wilf [13] showed that if G is K_{r+1} -free, then $\mu(G) \leq (1 - 1/r)n$. We strengthen these two results as follows.

Theorem 1 *If $G = G(n)$ is a K_{r+1} -free graph, then $\mu(G) < \mu(T_r(n))$ unless $G = T_r(n)$.*

Next, let G be a graph of order n , size m , and maximum degree $\Delta(G) = \Delta$. One of the best known facts about $\mu(G)$ is the inequality $\mu(G) \geq 2m/n$, due to Collatz and Sinogovitz [4]. In [11] we gave upper and lower bounds on $\mu(G) - 2m/n$ in terms of degree deviation. In turn, Cioabă and Gregory [3] showed that, if G is irregular and $n \geq 4$, then $\mu - 2m/n > 1/(n\Delta + 2n)$. In this note we give another proof of this bound and improve it for most graphs.

Call a graph *subregular* if $\Delta(G) - \delta(G) = 1$ and all but one vertices have the same degree.

Theorem 2 *If G is an irregular graph of order $n \geq 4$ and size m , then*

$$\mu(G) - 2m/n > 1/(2m + 2n) \quad (1)$$

unless G is subregular. If G is subregular with $\Delta(G) = \Delta$, then

$$\mu(G) - 2m/n > 1/(n\Delta + 2n). \quad (2)$$

Finally, write B_k for the graph consisting of k triangles sharing an edge, and let $0 \leq k \leq l \leq \Delta$. Shi and Song [12] showed that if $G = G(n)$ is a connected graph with $\Delta(G) = \Delta$, with no B_{k+1} and no $K_{2,l+1}$, then

$$\mu(G) \leq \left(k - l + \sqrt{(k - l)^2 + 4\Delta + 4l(n - 1)} \right) / 2. \quad (3)$$

We extend this result as follows.

Theorem 3 *Let $0 \leq k \leq l$. If $G = G(n)$ is a graph with $\Delta(G) = \Delta$, with no B_{k+1} and no $K_{2,l+1}$, then*

$$\mu(G) \leq \min \left\{ \Delta, \left(k - l + 1 + \sqrt{(k - l + 1)^2 + 4l(n - 1)} \right) / 2 \right\}. \quad (4)$$

If G is connected, equality holds if and only if one of the following conditions holds:

- (i) $\Delta^2 - \Delta(k - l + 1) \leq l(n - 1)$ and G is Δ -regular;
- (ii) $\Delta^2 - \Delta(k - l + 1) > l(n - 1)$ and every two vertices of G have k common neighbors if they are adjacent, and l common neighbors otherwise.

We note without a proof that (4) implies (3).

2 Proofs

Proof of Theorem 1 Write $k_r(G)$ for the number of r -cliques of G . The following result is given in [10]: if G is K_{r+1} -free graph, then

$$\mu^r(G) \leq \sum_{s=2}^r (s - 1) k_s(G) \mu^{r-s}(G). \quad (5)$$

According to a result of Zykov [14] (see also Erdős [5]), if the clique number of a graph G is r , then $k_s(G) < k_s(T_r(n))$ for every $2 \leq s \leq r$, unless $G = T_r(n)$. Assuming that $G \neq T_r(n)$, Zykov's theorem implies that $k_s(G) < k_s(T_r(n))$ for every $2 \leq s \leq r$. Hence, in view of (5), we have

$$\mu^r(G) < \sum_{s=2}^r (s-1) k_s(T_r(n)) \mu^{r-s}(G).$$

This implies that $\mu(G) < x$, where x is the largest root of the equation

$$x^r = \sum_{s=2}^r (s-1) k_s(T_r(n)) x^{r-s}. \quad (6)$$

It is known (see, e.g., [2], p. 74) that (6) is the characteristic equation of the Turán graph; so, $\mu(G) < x = \mu(T_r(n))$, completing the proof. $\square$

To simplify the proof of Theorem 2, we first prove inequality (1) for two special graphs.

Proposition 4 *Inequality (1) holds if G has $n-2$ vertices of degree $n-1$ and 2 vertices of degree $n-2$.*

Proof Clearly, G is the complete graph of order n with one edge removed. Using the theorem of Finck and Grohmann [6] (see also [2], Theorem 2.8), we find that

$$\mu(G) = \frac{n-3 + \sqrt{n^2 + 2n - 7}}{2}.$$

Hence, in view of $2m = n^2 - n - 2$, we obtain,

$$\begin{aligned} \mu(G) - \frac{2m}{n} &= \frac{\sqrt{n^2 + 2n - 7} - (n + 1 - \frac{4}{n})}{2} = \frac{4n - 8}{n^2 (\sqrt{n^2 + 2n - 7} + (n + 1 - \frac{4}{n}))} \\ &> \frac{4n - 8}{n^2 (n + 1 + (n + 1 - \frac{4}{n}))} \geq \frac{2n - 4}{n(n^2 + n - 2)} \geq \frac{1}{n^2 + n - 2} = \frac{1}{2m + 2n}, \end{aligned}$$

completing the proof. $\square$

Proposition 5 *Inequality (1) holds if G has $n-2$ vertices of degree $n-2$ and 2 vertices of degree $n-1$.*

Proof We easily deduce that n is even, say $n = 2k$, and that G is the complement of a $(k-1)$ -matching. Using the theorem of Finck and Grohmann, we find that

$$\mu(G) = \frac{n-3 + \sqrt{n^2 - 2n + 9}}{2}.$$

Hence, in view of $2m = n^2 - 2n + 2$, we obtain,

$$\begin{aligned}\mu(G) - \frac{2m}{n} &= \frac{\sqrt{n^2 - 2n + 9} - (n - 1 + \frac{4}{n})}{2} = \frac{4n - 8}{n^2 (\sqrt{n^2 - 2n + 9} + (n - 1 + \frac{4}{n}))} \\ &> \frac{4n - 8}{n^2 (n + 1 + (n - 1 + \frac{4}{n}))} = \frac{2n - 4}{n(n^2 + 2)} \geq \frac{1}{n^2 + 2} = \frac{1}{2m + 2n},\end{aligned}$$

completing the proof. $\square$

Proof of Theorem 2 Set $V = V(G)$, $\mu = \mu(G)$, and $\delta = \delta(G)$. Assume first that G is not subregular.

Proof of inequality (1)

Our proof is based on Hofmeister's inequality [9]: $\mu^2 \geq (1/n) \sum_{u \in V} d^2(u)$.

Case: $\Delta - \delta \geq 2$

In this case we easily see that

$$\sum_{u \in V} \left(d(u) - \frac{2m}{n} \right)^2 \geq 2 > \frac{2m}{m+n} + \frac{n}{4(m+n)^2},$$

and so,

$$\mu \geq \sqrt{\frac{1}{n} \sum_{u \in V} d^2(u)} = \sqrt{\frac{1}{n} \sum_{u \in V} \left(d(u) - \frac{2m}{n} \right)^2 + \frac{4m^2}{n^2}} > \frac{2m}{n} + \frac{1}{2m + 2n},$$

as claimed. Thus, hereafter we shall assume that $\Delta - \delta = 1$.

Case: $\Delta - \delta = 1$

Letting k be the number of vertices of degree $\Delta = \delta + 1$, we have $2m/n = \delta + k/n$, and so,

$$\frac{1}{n} \sum_{u \in V} \left(d(u) - \frac{2m}{n} \right)^2 = \frac{n-k}{n} \left(\frac{k}{n} \right)^2 + \frac{k}{n} \left(\frac{n-k}{n} \right)^2 = \frac{k(n-k)}{n^2}.$$

Hence, if

$$\frac{k(n-k)}{n^2} > \frac{2m}{n(m+n)} + \frac{1}{4(m+n)^2}, \quad (7)$$

then inequality (1) follows as above. Assume for contradiction that (7) fails.

Suppose first that either $k = 2$ or $n - k = 2$. Since (7) fails, we see that

$$\begin{aligned}2 - \frac{4}{n} &= \frac{(n-2)2}{n} \leq \frac{k(n-k)}{n} \leq \frac{2m}{m+n} + \frac{n}{4(m+n)^2} \\ &= 2 - \frac{2n}{m+n} + \frac{n}{4(m+n)^2}.\end{aligned} \quad (8)$$

In view of Propositions 4 and 5, we may assume that $\delta \leq n - 3$, and so

$$2m = \delta n + k \leq \delta n + n - 2 \leq n^2 - 2n - 2.$$

Noting that (8) increases in m , we obtain

$$-\frac{4}{n^2} \leq -\frac{4}{n^2 - 2} + \frac{1}{(n^2 - 2)^2},$$

a contradiction for $n \geq 4$.

Finally, let $k \geq 3$ and $n - k \geq 3$; thus, $n \geq 6$. We have

$$2m = \delta n + k \leq \delta n + n - 3 \leq (n - 2)n + n - 3.$$

By assumption inequality (7) fails; hence,

$$3 - \frac{9}{n} \leq \frac{(n - k)k}{n} \leq 2 - \frac{2n}{m + n} + \frac{n}{4(m + n)^2} \leq 2 - \frac{4n}{n^2 + n - 3} + \frac{n}{(n^2 + n - 3)^2}.$$

This inequality is a contradiction for $n \geq 6$, completing the proof of (7).

Proof of inequality (2) when G is subregular

Since G is subregular, it has either a single vertex of degree Δ or a single vertex of degree δ . Clearly, $\delta \geq 1$, and so $m > n/2$.

Case: G has a single vertex of degree Δ

Setting $\Delta = k + 1$ and

$$c = \frac{nk + 1}{n} + \frac{1}{n(k + 3)} = k + \frac{k + 4}{n(k + 3)},$$

in view of $2m = nk + 1$, inequality (2) amounts to $\mu > c(G)$.

Select a vertex $u \in V$ with $d(u) = k + 1$; partition V as $V = \{u\} \cup V \setminus \{u\}$ and let B be the quotient matrix of this partition (see, e.g. [8], Ch. 9), i.e.,

$$B = \begin{pmatrix} 0 & \frac{k+1}{n-1} \\ k+1 & k - \frac{k+1}{n-1} \end{pmatrix}.$$

Writing $P(x)$ for the characteristic polynomial of B and observing that $k \leq n - 2$, we have

$$\begin{aligned} P(c) &= \left(k + \frac{k+4}{n(k+3)}\right) \left(k + \frac{k+4}{n(k+3)} - \left(k - \frac{k+1}{n-1}\right)\right) - \frac{(k+1)^2}{n-1} \\ &= k \frac{k+4}{n(k+3)} + \frac{1}{n^2} \left(\frac{k+4}{k+3}\right)^2 + \left(\frac{k+4}{n(k+3)}\right) \frac{k+1}{n-1} - \frac{k+1}{n-1} \\ &= -\frac{3}{n(k+3)} + \frac{1}{n^2} \left(\frac{k+4}{k+3}\right)^2 + \frac{(k+1)}{n(n-1)(k+3)} \\ &= \frac{1}{n^2(k+3)} \left(-3n + 2k + 6 + \frac{1}{k+3} + \frac{k+1}{n-1}\right) \\ &\leq \frac{1}{n^2(k+3)} \left(-3n + 2(n-2) + 6 + \frac{1}{4} + 1\right) < 0. \end{aligned}$$

By interlacing, $P(\mu) \geq 0 > P(c)$, and so $\mu > c$, completing the proof of (2) in this case.

Case: G has a single vertex of degree δ

Setting $\Delta = k$ and

$$c = \frac{nk - 1}{n} + \frac{1}{n(k+3)} = k - \frac{k+1}{n(k+2)},$$

in view of $2m = nk - 1$, inequality (2) amounts to $\mu > c$.

Select $u \in V$ with $d(u) = k - 1$; partition V as $V = \{u\} \cup V \setminus \{u\}$ and let B be the quotient matrix of this partition, i.e.,

$$B = \begin{pmatrix} 0 & \frac{k-1}{n-1} \\ k-1 & k - \frac{k-1}{n-1} \end{pmatrix}.$$

Writing $P(x)$ for the characteristic polynomial of B and observing that $k \leq n - 2$, we have

$$\begin{aligned} P(c) &= \left(k - \frac{k+1}{n(k+2)}\right) \left(k - \frac{k+1}{n(k+2)} - \left(k - \frac{k-1}{n-1}\right)\right) - \frac{(k-1)^2}{n-1} \\ &= -\frac{k(k+1)}{n(k+2)} + \frac{1}{n^2} \left(\frac{k+1}{k+2}\right)^2 + \frac{k-1}{n(n-1)(k+2)} + \frac{k-1}{n} \\ &= -\frac{2}{n(k+2)} + \frac{1}{n^2} \left(\frac{k+1}{k+2}\right)^2 + \frac{k-1}{n(n-1)(k+2)} \\ &= \frac{1}{n^2(k+2)} \left(-2n + 2k + 1 + \frac{1}{k+2} + \frac{k-1}{n-1}\right) \\ &< \frac{1}{n^2(k+2)} \left(-2n + 2(n-2) + 1 + \frac{1}{1+2} + 1\right) < 0. \end{aligned}$$

By interlacing, $P(\mu) \geq 0 > P(c)$, completing the proof of (2). $\square$

Proof of Theorem 3 Set $V = V(G)$ and $\mu = \mu(G)$; given $u \in V$, write $\Gamma(u)$ for the set of neighbors of u . Select $u \in V$; let $A = \Gamma(u)$, $B = V \setminus (\Gamma(u) \cup \{u\})$, and $e(A, B)$ be the number of $A - B$ edges. Since G contains no B_{k+1} and no $K_{2,l+1}$, we see that

$$\sum_{v \in A} (d(v) - k - 1) \leq \sum_{v \in A} |\Gamma(v) \cap B| = e(A, B) = \sum_{v \in A} |\Gamma(v) \cap A| \leq (n - d(u) - 1)l. \quad (9)$$

Letting A be the adjacency matrix of G , note that the u th row sum of the matrix

$$C = A^2 - (k+1-l)A - (n-1)lI_n$$

is equal to

$$\sum_{v \in A} (d(v) - k - 1) - (n-1-d(u))l;$$

consequently, all row sums C are nonpositive. Letting $\mathbf{x} = (x_1, \dots, x_n)$ be an eigenvector of A to μ , we see that the value

$$\lambda = \mu^2 - (k + 1 - l)\mu - (n - 1)l$$

is an eigenvalue of C with eigenvector $\mathbf{x}$. Therefore, $\lambda \leq 0$, and so,

$$\mu \leq \left(k - l + 1 + \sqrt{(k - l + 1)^2 + 4l(n - 1)} \right) / 2,$$

completing the proof of inequality (4).

Let equality hold in (4) and G be connected; thus, the eigenvector $\mathbf{x} = (x_1, \dots, x_n)$ to μ is positive. We shall prove the necessity of conditions (i) and (ii). If

$$\mu = \Delta \leq \left(k - l + 1 + \sqrt{(k - l + 1)^2 + 4l(n - 1)} \right) / 2,$$

then $\Delta^2 - \Delta(k - l + 1) \leq l(n - 1)$ and G is Δ -regular.

On the other hand, if

$$\mu = \left(k - l + 1 + \sqrt{(k - l + 1)^2 + 4l(n - 1)} \right) / 2 < \Delta,$$

then $\Delta^2 - \Delta(k - l + 1) > l(n - 1)$ and $\lambda = 0$. Scaling $\mathbf{x}$ so that $x_1 + \dots + x_n = 1$, we see that λ is a convex combination of the row sums of C which are nonpositive; thus, all row sums of C are 0. Since equality holds in (9) for every $u \in [n]$, every two vertices have exactly k common neighbors if they are adjacent, and exactly l common neighbors otherwise. This completes the proof. $\square$

3 Concluding remarks

Finding tight bounds on the spectral radius of subregular graphs is a challenging problem. Specifically, we cannot determine for which subregular graphs G one has

$$\mu(G) > \frac{2m}{n} + \frac{1}{2m + 2n}.$$

Note that strongly regular graphs satisfy condition (ii) for equality in (4), but irregular graphs can satisfy this condition as well, e.g., the star $K_{1,n-1}$ and the friendship graph.

Finally, setting $l = \Delta$ or $k = 0$, Theorem 3 implies assertions that strengthen Corollaries 1 and 2 of [12].

References

- [1] B. Bollobás, *Modern Graph Theory*, Graduate Texts in Mathematics, **184**, Springer-Verlag, New York (1998), xiv+394 pp.
- [2] D. Cvetković, M. Doob, and H. Sachs, *Spectra of Graphs*, VEB Deutscher Verlag der Wissenschaften, Berlin, 1980, 368 pp.
- [3] S. M. Cioabă, D. Gregory, Large matchings from eigenvalues, to appear in *Linear Algebra Appl.*
- [4] L. Collatz, U. Sinogowitz, Spektren endlicher Grafen, *Abh. Math. Sem. Univ. Hamburg* **21** (1957), 63-77.
- [5] P. Erdős, On the number of complete subgraphs contained in certain graphs, *Publ. Math. Inst. Hung. Acad. Sci.* **VII**, Ser. A3 (1962), 459-464p.
- [6] H.J.Finck, G. Grohmann, Vollständiges Produkt, chromatische Zahl und charakteristisches Polynom regulärer Graphen. I. (German) *Wiss. Z. Techn. Hochsch. Ilmenau* **11** (1965) 1-3.
- [7] L. Feng, Q. Li, X.-D. Zhang, Spectral radii of graphs with given chromatic number, *Appl. Math. Letters*, **20** (2007), 158-162.
- [8] C. Godsil, G. Royle, *Algebraic graph theory*, Graduate Texts in Mathematics, **207**, Springer-Verlag, New York (2001), xx+439 pp.
- [9] M. Hofmeister, Spectral radius and degree sequence, *Math. Nachr.* **139** (1988), 37-44.
- [10] V. Nikiforov, Some inequalities for the largest eigenvalue of a graph. *Combin. Probab. Comput.* **11** (2002), 179-189.
- [11] V. Nikiforov, Eigenvalues and degree deviation in graphs, *Linear Algebra Appl.* **414** (2006), 347-360.
- [12] L. Shi, Z. Song, Upper bounds on the spectral radius of book-free and/or $K_{2,l+1}$ graphs, *Linear Algebra Appl.* **420** (2007), 526-529.
- [13] H. Wilf, Spectral bounds for the clique and independence numbers of graphs, *J. Combin. Theory Ser. B* **40** (1986), 113-117.
- [14] A. A. Zykov, On some properties of linear complexes (in Russian), *Mat. Sbornik N.S.* **24**(66), (1949), 163-188.